

Questioning the evidence for a Central Asian domestication origin of dogs

Guo-Dong Wang^{a,b}, Min-Sheng Peng^a, He-Chuan Yang^a, Peter Savolainen^c, and Ya-Ping Zhang^{a,d,1}

A recent study by Shannon et al. (1) proposes a Central Asian domestication origin of dogs, based upon the finding that Central Asia had the lowest short-range linkage disequilibrium (LD) among village dogs across the globe. A reanalysis of their data, however, suggests that these conclusions may require revision.

First, Shannon et al.'s (1) definition of "Central Asia" is questionable. Mongolia and Nepal [the two regions assigned to Central Asia by Shannon et al. (1)] are not associated with the common definition of Central Asia that includes the region bounded by the Caspian Sea, Russia, China, and Afghanistan, but are located in East Asia and South Asia, respectively. Although Mongolia is occasionally considered a part of Central Asia (e.g., by the United Nations Educational, Scientific, and Cultural Organization), Nepal is not, but is considered a part of South Asia according to all modern definitions (Fig. 1A). There is no expectation that the dogs in these countries represent a single, interbreeding population. Using the data from Shannon et al. (1), we recalculated the LD of Mongolia and Nepal separately, finding different LD patterns in the two countries (Fig. 1B). The LD is lower for Nepal than for the combined population, at both short inter-SNP distances (<0.0005 cM) and intermediate distances (0.01–0.05 cM), and higher for Mongolia, suggesting that the assumption of a single population is not correct.

Most important, although several studies have argued that domestic dogs originated in southern China (2), no dogs from this region were included in Shannon et al.'s analysis (1), neglecting available published data (3, 4). We therefore included published data from Ying Jiang, southwestern China (3, 4) (Fig. 1A), in the LD analysis. The results show that this South Chinese population, and not Central Asia or Nepal, had the lowest LD of all populations at both short and intermediate inter-SNP distances (Fig. 1B). Thus,

although we question this statistic as a sole indicator of dog geographical origin, we here show that if low LD at short inter-SNP distances is used to identify the geographical origin of the dog, the available data suggest southern China rather than Central Asia or Nepal.

Last, Shannon et al. (1) correctly note that East Asia contains the largest number of mitochondrial DNA haplogroups worldwide, but claim that East Asia contains fewer Y haplogroups than most other regions, by lacking haplogroup 8 (H8). However, this assertion does not take published data into account. H8 was previously found in an East Asian dog, actually the "type specimen" in which H8 was first identified (5), implying that East Asia contains at least the same number of Y haplogroups as all other geographical regions.

These issues, related to geographical partitions and regional sampling bias, suggest that the conclusions reported by Shannon et al. (1) may require revision. It is important that future studies of dog origins incorporate samples from across the Old World, including southern China and Southeast Asia, in broad collaborations, and that data quality can be normalized across different data types.

Acknowledgments

We thank Greger Larson and Laurent Frantz at the University of Oxford for critical input. This work was supported by grants from the NSFC (Grant 91531303), the 973 programs (Grants 2013CB835200 and 2013CB835202), and the Breakthrough Project of Strategic Priority Program of the Chinese Academy of Sciences (Grant XDB13000000). G.-D.W. is supported by a grant from the China Scholarship Council (Grant 201504910368). G.-D. W. and M.-S.P. are supported by the Youth Innovation Promotion Association, Chinese Academy of Sciences.

^aState Key Laboratory of Genetic Resources and Evolution & Yunnan Laboratory of Molecular Biology of Domestic Animals & Germplasm Bank of Wild Species, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China; ^bPalaeogenomics & Bio-Archaeology Research Network, Department of Archaeology, Research Laboratory for Archaeology and the History of Art, University of Oxford, Oxford OX1 3QY, United Kingdom; ^cDepartment of Gene Technology, KTH-Royal Institute of Technology, Science for Life Laboratory, 17165 Solna, Sweden; and ^dLaboratory for Conservation and Utilization of Bio-Resources & Key Laboratory for Microbial Resources of the Ministry of Education, Yunnan University, Kunming 650091, China

Author contributions: G.-D.W. and Y.-P.Z. designed research; M.-S.P. and H.-C.Y. analyzed data; and G.-D.W., M.-S.P., and P.S. wrote the paper. The authors declare no conflict of interest.

¹To whom correspondence should be addressed. Email: zhangyp@mail.kiz.ac.cn.




Fig. 1. (A) Map showing the location of Mongolia and Nepal, and Ying Jiang in southern China. **(B)** LD decay curves for village dogs worldwide based on the data from Shannon et al. (1), and the Ying Jiang village dogs. The dashed lines represent the populations from Mongolia (gray) and Nepal (blue), respectively. The grass green line represents the population from Ying Jiang, southern China.

- 1 Shannon LM, et al. (2015) Genetic structure in village dogs reveals a Central Asian domestication origin. *Proc Natl Acad Sci USA* 112(44):13639–13644.
- 2 Wang GD, et al. (2016) Out of southern East Asia: The natural history of domestic dogs across the world. *Cell Res* 26(1):21–33.
- 3 Bai B, et al. (2015) DoGSD: The dog and wolf genome SNP database. *Nucleic Acids Res* 43(Database issue):D777–D783.
- 4 Gou X, et al. (2014) Whole-genome sequencing of six dog breeds from continuous altitudes reveals adaptation to high-altitude hypoxia. *Genome Res* 24(8): 1308–1315.
- 5 Natanaelsson C, et al. (2006) Dog Y chromosomal DNA sequence: Identification, sequencing and SNP discovery. *BMC Genet* 7:45.